



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES
(untabulated).

AZORES—*St. Michaels.*—Month of May, 1910. Population, 131,183. Total number of deaths from all causes 276, including tuberculosis 7.

CANADA—*Vancouver.*—Month of July, 1910. Population, 78,900. Total number of deaths from all causes 71, including measles 1 and tuberculosis 9.

Yarmouth.—Month of July, 1910. Population, 6,700. Total number of deaths from all causes 9, including diphtheria 1 and tuberculosis 1.

CHILE—*Punta Arenas.*—Month of April, 1910. Population, 12,000. Total number of deaths from all causes 32, including tuberculosis 4.

EGYPT—*Alexandria.*—Month of June, 1910. Population, 720,000. Total number of deaths from all causes not reported. Deaths from contagious diseases include typhus fever 5, typhoid fever 9, smallpox 3, measles 38, scarlet fever 1, diphtheria 4, plague 3, tuberculosis 63.

FRANCE—*Nice.*—Month of June, 1910. Population, 150,881. Total number of deaths from all causes, 204, including typhoid fever 2, measles 1, diphtheria 1, tuberculosis 31.

GREAT BRITAIN.—Week ended July 23, 1910.

England and Wales.—The deaths registered in 77 great towns correspond to an annual rate of 11.1 per 1,000 of the population, which is estimated at 16,940,895.

Ireland.—The deaths registered in 21 principal town districts correspond to an annual rate of 16.3 per 1,000 of the population, which is estimated at 1,151,790. The lowest rate was recorded at Dundalk, viz, 4, and the highest at Kilkenny, viz, 24.6.

Scotland.—The deaths registered in 8 principal towns correspond to an annual rate of 12 per 1,000 of the population, which is estimated at 1,865,571. The lowest rate was recorded at Greenock, viz, 7.8, and the highest at Perth, viz, 19.6. The total number of deaths from all causes was 434, including typhoid fever 4, measles 5, scarlet fever 6, diphtheria 8.

MAURITIUS—*Port Louis.*—Month of April, 1910. Population, 373,071. Total number of deaths from all causes 998, including plague 3 and tuberculosis 57.

RUSSIA—*Riga.*—Month of May, 1910. Population, 355,000. Total number of deaths from all causes 719, including typhus fever 8, typhoid fever 7, smallpox 26, measles 6, scarlet fever 52, diphtheria 15, tuberculosis 84.

SPAIN—*Huelva.*—Month of June, 1910. Population 24,000. Total number of deaths from all causes 80, including measles 1, scarlet fever 1, diphtheria 2, tuberculosis 12.

URUGUAY—*Montevideo*. — Month of May, 1910. Population, 321,224. Total number of deaths from all causes 528, including typhoid fever 6, smallpox 102, scarlet fever 1, diphtheria 3, tuberculosis 55.

By authority of the Secretary of the Treasury:

WALTER WYMAN,

Surgeon-General,

United States Public Health and Marine-Hospital Service.